

MATRICES

1.- Calculate the following determinants:

$$a) \begin{vmatrix} 3 & 2 & 5 \\ 2 & 1 & 4 \\ 3 & 1 & 6 \end{vmatrix}$$

$$b) \begin{vmatrix} 2 & 1 & 3 \\ 4 & 2 & 5 \\ 6 & 0 & -2 \end{vmatrix}$$

$$c) \begin{vmatrix} 3 & 1 & 5 & 0 \\ 5 & 4 & 6 & 3 \\ 1 & 3 & 2 & 1 \\ 6 & 7 & 5 & 4 \end{vmatrix}$$

$$d) \begin{vmatrix} 7 & 6 & 8 & 5 \\ 6 & 7 & 10 & 6 \\ 7 & 8 & 8 & 9 \\ 8 & 7 & 9 & 6 \end{vmatrix}$$

$$e) \begin{vmatrix} 1 & 3 & 2 & 1 \\ 3 & 5 & 3 & 2 \\ 3 & 6 & 2 & 2 \\ 6 & 4 & 5 & 3 \end{vmatrix}$$

$$f) \begin{vmatrix} -1 & 1 & 1 & 1 \\ 1 & -1 & 1 & 1 \\ 1 & 1 & -1 & 1 \\ 1 & 1 & 1 & -1 \end{vmatrix}$$

$$g) \begin{vmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & 1 \\ 1 & 1 & -1 & 1 \\ 1 & 1 & 1 & -1 \end{vmatrix}$$

$$h) \begin{vmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{vmatrix}$$

$$i) \begin{vmatrix} 3 & 1 & 1 & 1 \\ 1 & 3 & 1 & 1 \\ 1 & 1 & 3 & 1 \\ 1 & 1 & 1 & 3 \end{vmatrix}$$

$$j) \begin{vmatrix} 3 & 4 & 0 & 0 \\ 4 & 2 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{vmatrix}$$

$$k) \begin{vmatrix} 3 & 4 & 1 & 2 \\ 0 & 3 & 1 & 5 \\ 0 & 0 & 2 & 7 \\ 0 & 0 & 0 & 4 \end{vmatrix}$$

$$l) \begin{vmatrix} 1 & 2 & 1 & 2 & 1 \\ 0 & 0 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 2 \\ 1 & 2 & 2 & 1 & 1 \end{vmatrix}$$

$$m) \begin{vmatrix} 1 & 2 & 3 & 4 & 5 \\ -1 & 0 & 3 & 4 & 5 \\ -1 & -2 & 0 & 4 & 5 \\ -1 & -2 & -3 & 0 & 5 \\ -1 & -2 & -3 & -4 & 0 \end{vmatrix}$$

$$n) \begin{vmatrix} 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 2 & 1 \\ 0 & 0 & 3 & 2 & 1 \\ 0 & 4 & 3 & 2 & 1 \\ 5 & 4 & 3 & 2 & 1 \end{vmatrix}$$

$$ñ) \begin{vmatrix} 0 & a & b \\ a & 0 & c \\ b & c & 0 \end{vmatrix}$$

$$o) \begin{vmatrix} a & b & b \\ b & a & b \\ b & b & a \end{vmatrix}$$

$$p) \begin{vmatrix} x & 1 & 1 \\ 1 & x & 1 \\ 1 & 1 & x \end{vmatrix}$$

$$q) \begin{vmatrix} x+a & b & c \\ a & x+b & c \\ a & b & x+c \end{vmatrix}$$

$$r) \begin{vmatrix} a & 3 & 0 & 5 \\ 0 & b & 0 & 2 \\ 1 & 2 & c & 3 \\ 0 & 0 & 0 & d \end{vmatrix}$$

$$s) \begin{vmatrix} 1 & 1 & 1 & 1 & . & . & 1 \\ -1 & a & 1 & 1 & . & . & 1 \\ -1 & -1 & a & 1 & . & . & 1 \\ . & . & . & . & . & . & . \\ -1 & -1 & -1 & -1 & . & . & a \end{vmatrix}$$

$$t) \begin{vmatrix} 1 & 2 & 3 & . & . & n-1 \\ 2 & 3 & 4 & . & . & n \\ 3 & 4 & 5 & . & . & n+1 \\ . & . & . & . & . & . \\ n-2 & n-1 & n & . & . & 2n-2 \\ n-1 & n & n+1 & . & . & 2n-1 \end{vmatrix}$$

2.- Solve the equation $\begin{vmatrix} 4 & x & 6 \\ 5 & 7 & 12 \\ 3 & -1 & x \end{vmatrix} = 0$.

3.- Show that:

a) $\begin{vmatrix} x & y & x+y \\ y & x+y & x \\ x+y & x & y \end{vmatrix} = -2(x^3 + y^3)$ b) $\begin{vmatrix} 1+x & 1 & 1 & 1 \\ 1 & 1-x & 1 & 1 \\ 1 & 1 & 1+z & 1 \\ 1 & 1 & 1 & 1-z \end{vmatrix} = x^2 z^2$

4.- Let $A = \begin{pmatrix} 2 & 4 & 1 \\ 6 & -3 & 2 \\ 4 & 1 & 3 \end{pmatrix}$. Calculate the determinant $|A - \lambda I_3|$, with $\lambda \in \mathbb{R}$.

5.- Knowing that $\begin{vmatrix} x & y & z \\ 3 & 0 & 2 \\ 1 & 1 & 1 \end{vmatrix} = 1$, obtain the value of $\begin{vmatrix} x & y & z \\ 3x+3 & 3y & 3z+2 \\ x+4 & y+4 & z+4 \end{vmatrix}$.

6.- Let $A, B \in M_4$ with $|A|=3$, $|B|=-2$. Calculate:

a) $|2A|$.

b) $|\frac{1}{2}B|$.

c) $|BA^t|$.

d) $|(BA)^t|$.

e) $|(B^t A^t B)^t|$.

7.- Calculate the rank of the following matrices:

$$A = \begin{pmatrix} 2 & -2 & -4 \\ -1 & 3 & 4 \\ 1 & -2 & -3 \end{pmatrix}$$

$$B = \begin{pmatrix} 2 & 0 & 3 \\ 4 & 1 & 0 \\ 10 & 1 & 5 \end{pmatrix}$$

$$C = \begin{pmatrix} 2 & -1 & 0 & 1 & 3 \\ 1 & 0 & -1 & 2 & 3 \\ 3 & -1 & -1 & 3 & 6 \\ 5 & -2 & -1 & 4 & 9 \end{pmatrix}$$

$$D = \begin{pmatrix} -2 & -4 & 3 & 4 & 1 & 8 \\ 1 & 2 & 0 & -2 & 1 & -1 \\ 3 & 6 & 3 & -6 & 6 & 3 \\ 2 & 4 & 2 & -4 & 4 & 2 \end{pmatrix}$$

$$E = \begin{pmatrix} 1 & -2 & 1 & 0 & 2 \\ 0 & 1 & 2 & -1 & 1 \\ 1 & 2 & 0 & -2 & 1 \\ 2 & 1 & 3 & -3 & 4 \\ 3 & -2 & 2 & -2 & 5 \end{pmatrix}$$

$$F = \begin{pmatrix} 1 & 2 & 3 \\ -1 & 1 & 1 \\ 3 & 3 & 5 \\ 0 & 3 & 4 \end{pmatrix}$$

$$G = \begin{pmatrix} 1 & 1 & -2 & 1 \\ 2 & 0 & 1 & 3 \\ -1 & 1 & 2 & 1 \\ 3 & 2 & 1 & 2 \end{pmatrix} \quad H = \begin{pmatrix} 0 & 2 & 1 & -1 & 2 \\ 1 & 0 & -1 & 3 & 2 \\ 0 & 4 & 2 & -2 & 4 \end{pmatrix} \quad I = \begin{pmatrix} 1 & 2 \\ 0 & 1 \\ 1 & 0 \end{pmatrix}$$

8.- Calculate the rank of the following matrices depending on the values of the real parameter a :

$$A = \begin{pmatrix} 1 & 1 & 0 & 1 \\ 3 & 2 & -1 & 3 \\ a & 3 & -2 & 0 \\ -1 & 0 & -4 & 3 \end{pmatrix} \quad B = \begin{pmatrix} 1 & 1 & -1 & 1 \\ a & 1 & 1 & 1 \\ 1 & -1 & 3 & 0 \\ 4 & 2 & 0 & a \end{pmatrix} \quad C = \begin{pmatrix} 1 & -1 & 2 \\ 2 & 1 & 3 \\ 5 & 1 & a \end{pmatrix}$$

9.- Study if the following matrices are invertible. If affirmative, calculate the inverse matrix.

$$A = \begin{pmatrix} 0 & 1 & 2 \\ 2 & 3 & 0 \\ 4 & 5 & 7 \end{pmatrix} \quad B = \begin{pmatrix} 1 & 2 & 0 \\ 0 & 2 & 0 \\ 0 & 2 & 1 \end{pmatrix} \quad C = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix} \quad D = \begin{pmatrix} 0 & 3 & 1 \\ -3 & 0 & -2 \\ -1 & 2 & 0 \end{pmatrix}$$

$$E = \begin{pmatrix} 1 & -1 & 3 \\ 0 & 4 & -2 \\ -2 & 6 & -8 \end{pmatrix} \quad F = \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{pmatrix} \quad G = \begin{pmatrix} 2 & -1 & 1 & 1 \\ 0 & 0 & 1 & 0 \\ 2 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

10.- Study the existence of the inverse matrix depending on the values of $m \in \mathbb{R}$. Calculate the inverse matrix in the invertible cases.

$$A = \begin{pmatrix} m & -1 & 0 \\ 1 & m-1 & 1 \\ 1 & 0 & 1 \end{pmatrix} \quad B = \begin{pmatrix} m & 1 & -1 \\ 1 & 2 & 1 \\ 1 & 3 & -1 \end{pmatrix} \quad C = \begin{pmatrix} m & 1 & 1 \\ 1 & m & 1 \\ 1 & 1 & m \end{pmatrix}$$

11.- Study for what values of the parameter $m \in \mathbb{R}$, the following matrices do not have inverse.

$$A = \begin{pmatrix} 1-m & 2 \\ 3 & 2-m \end{pmatrix} \quad B = \begin{pmatrix} -m & 5 \\ 2 & 3-m \end{pmatrix} \quad C = \begin{pmatrix} 2-m & 3 & 1 \\ 1 & 1-m & 4 \\ 0 & 1 & 1-m \end{pmatrix}$$

$$D = \begin{pmatrix} m & 9 & 4 \\ 4 & m & -1 \\ 7 & 7 & 7 \end{pmatrix} \quad E = \begin{pmatrix} 1 & 1 & 1 \\ m & 1 & 0 \\ -1 & 3 & m-1 \end{pmatrix}$$

12.- Given the matrices $P = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix}$ and $B = \begin{pmatrix} 1 & 2 & -1 \\ 0 & -3 & 1 \\ 2 & 0 & 1 \end{pmatrix}$, calculate the matrix A that

verifies $P^{-1}AP = B$.

13.- Solve, using Cramer's Rule:

$$\begin{array}{lll} \text{a)} \quad \begin{vmatrix} x+y-z=3 \\ 5x-y+2z=5 \\ -3x+3y-4z=1 \end{vmatrix} & \text{b)} \quad \begin{vmatrix} x+y-z=1 \\ x+2y+z=2 \\ x+3y-z=0 \end{vmatrix} & \text{c)} \quad \begin{vmatrix} x+y-2z+t+3s=1 \\ 2x-y+2z+2t+6s=2 \\ 3x+2y-4z-3t-9s=3 \end{vmatrix} \end{array}$$

14.- Discuss the following systems of linear equations as the real parameters (m , a and b) varies, and solve them using Cramer's Rule:

$$\begin{array}{lll} \text{a)} \quad \begin{vmatrix} mx+y-z=1 \\ x+2y+z=2 \\ x+3y-z=0 \end{vmatrix} & \text{b)} \quad \begin{vmatrix} mx+y+z=m \\ x+my+z=1 \\ x+y+mz=1 \end{vmatrix} & \text{c)} \quad \begin{vmatrix} x+y+z=m \\ x+(1+m)y+z=2m \\ x+y+z=4 \end{vmatrix} \\ \text{d)} \quad \begin{vmatrix} x+my-z=0 \\ 2x-3y-2z=0 \\ x+2y+z=0 \end{vmatrix} & \text{e)} \quad \begin{vmatrix} mx+y+3z=3 \\ x-y-z=0 \\ 5x-3y-2z=6 \end{vmatrix} & \text{f)} \quad \begin{vmatrix} x-2y+z=-1 \\ x+y+3z=4 \\ 5x-y+mz=10 \end{vmatrix} \\ \text{g)} \quad \begin{vmatrix} -x+2y-2z=0 \\ 2x-y+az=b \\ 2x-2y+3z=1+b \end{vmatrix} & \text{h)} \quad \begin{vmatrix} 2x+ay+z=7 \\ x+ay+z+t=b \\ x+2ay+t=-1 \\ bx+ay=b \end{vmatrix} & \end{array}$$

15.- Given the matrix $A = \begin{pmatrix} 1 & 1 & m \\ 3 & 2 & 4m \\ 2 & 1 & 3 \end{pmatrix}$ with $m \in \mathbb{R}$:

a) For what values of the parameter m , is the matrix A invertible?

b) Solve the linear system $AX = 0_3$ using the results of point a).

16.- Consider the matrices $A = \begin{pmatrix} 1 & a & 0 & 0 \\ a & 1 & 0 & 1-a \\ 0 & 0 & 1 & 0 \end{pmatrix}$, $B = \begin{pmatrix} a \\ a^2 \\ a^3 \end{pmatrix}$ and $X = \begin{pmatrix} x \\ y \\ z \\ t \end{pmatrix}$, with a being

a real parameter.

a) Discuss the system $AX = B$ as a varies.

b) In the compatible cases, calculate the solutions of $AX = B$.

17 .- In a market with perfect competition the functions of supply and demand of the goods are given by:

$$Q_{1d} = 10 - 2P_1 + 4P_2$$

$$Q_{2d} = 10 + 5P_1 - 3P_2$$

$$Q_{1s} = 20 + 3P_1 - 2P_2$$

$$Q_{2s} = 30 - 7P_1 + 5P_2$$

Where Q_{id} is the demanded quantity of good i , Q_{is} is the supplied quantity of good i and P_i is the price of the good i , for $i=1,2$. Calculate the prices for which the market is in equilibrium and the demanded and supplied quantities of each good in this situation.

18 .- The condition of equilibrium for the price of three goods in a given market is determined by the following equations:

$$11P_1 - P_2 - P_3 = 31$$

$$-P_1 + 6P_2 - 2P_3 = 26$$

$$-P_1 - 2P_2 + 7P_3 = 24$$

where P_1 , P_2 and P_3 are the prices of these three goods. Calculate the price of equilibrium for each good.

19.- Consider the vectors $u = (1, -3, 2)$ and $v = (2, -1, 1)$ of \mathbb{R}^3 :

- Write, if it is possible, the vectors $(1, 7, -4)$ and $(2, -5, 4)$ as a linear combination of u and v .
- For which values of x is the vector $(1, x, 5)$ a linear combination of u and v ?

20.- The vectors $v_1 = (1, 1, -1)$, $v_2 = (2, 1, 3)$ and $v_3 = (5, 2, 10)$ of \mathbb{R}^3 , are linearly independent? If not, find the relation of dependency.

21.- Given the vectors $u_1 = (2, -1, 0)$, $u_2 = (0, 1, -1)$ and $u_3 = (8, 3, 1)$ of \mathbb{R}^3 , study if they are linearly dependent or independent.

22.- Given the vectors $u_1 = (1, 0, -1, 0)$, $u_2 = (2, 0, 3, -1)$, $u_3 = (1, 1, -1, 1)$ and $u_4 = (2, 1, -2, 1)$ of \mathbb{R}^4 , study if they are linearly independent. If not, find the relation of dependency.

23.- Given the vectors of \mathbb{R}^4 $v_1 = (1, 1, 0, m)$, $v_2 = (3, -1, n, -1)$ and $v_3 = (-3, 5, m, -4)$, determine the values of the parameters m and n so that the three vectors be linearly dependent.

24.- Sean $u = (-1, 0, 0)$, $v = (1, 1, 0)$ and $w = (-1, 1, -1)$ vectors of \mathbb{R}^3 :

- Show that $\{u, v, w\}$ is a basis of \mathbb{R}^3 .
- Find the coordinates with respect to this basis of the vector whose coordinates with respect to the canonical basis are 1, 0, 2.
- Find the coordinates concerning the canonical basis of the vector $a = 3u - v + 5w$

25.- Let $A \in M_n$, $\lambda \in \mathbb{R}$ an eigenvalue of A and $\mathbf{x} \in \mathbb{R}^n$ an eigenvector of A associated to λ

Prove that:

- $\alpha\lambda$ is an eigenvalue of the matrix αA for any $\alpha \in \mathbb{R}$ and \mathbf{x} is an eigenvector of αA associated to $\alpha\lambda$.
- λ^p is an eigenvalue of A^p and \mathbf{x} is an eigenvector of A^p associated to λ^p , with $p \in \mathbb{N}$.
- $|A| = 0 \Leftrightarrow \lambda = 0$ is an eigenvalue of A .
- If A invertible then $\lambda \neq 0$. Moreover, λ^{-1} is an eigenvalue of A^{-1} and \mathbf{x} is an eigenvector of A^{-1} associated to λ^{-1} .

26.- Let $A, B \in M_n$ be similar matrices. Prove that:

- $|A| = |B|$.
- A^p is similar to B^p for any $p \in \mathbb{N}$.
- If A is invertible then B is invertible and A^{-1} is similar to B^{-1} .

27.- Let $A \in M_n$. Prove that A and A^t have the same characteristic polynomial.

28.- Given the matrix $A = \begin{pmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 3 & 0 & 2 \end{pmatrix}$ answer the following questions:

- Study if 3 is eigenvalue of A or not.
- Are the vectors (1,1,1) and (0,0,1) eigenvectors of A ? If in the affirmative, find the associated eigenvalues.

29.- Given the matrix $A = \begin{pmatrix} -1 & 0 & 0 \\ 1 & 2 & 0 \\ 0 & -3 & 1 \end{pmatrix}$ answer the following questions:

- Study if the vector $(-1, \frac{1}{3}, \frac{1}{2})$ is an eigenvector or not of the matrix A . If in the affirmative, determine the associated eigenvalue.
- The same question for the vector $(-1, 0, 1)$.

30.- For each one of the following matrices reason out if it is diagonalizable or not. Besides:

- If in the affirmative, give a diagonal matrix D and an invertible matrix P such that $D = P^{-1}AP$.
- If in the negative, calculate the eigenvalues and the eigenvectors associated to each eigenvalue.

$$A_1 = \begin{pmatrix} 0 & 3 & 0 \\ -1 & 4 & 0 \\ 0 & 2 & -2 \end{pmatrix}$$

$$A_2 = \begin{pmatrix} 2 & 2 & 1 \\ -2 & 1 & 2 \\ 1 & 2 & 2 \end{pmatrix}$$

$$A_3 = \begin{pmatrix} 1 & -3 & 3 \\ 3 & -5 & 3 \\ 6 & -6 & 4 \end{pmatrix}$$

$$A_4 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}$$

$$A_5 = \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ -1 & 1 & 0 \end{pmatrix}$$

$$A_6 = \begin{pmatrix} 4 & -4 & 6 \\ 3 & -4 & 6 \\ 1 & -2 & 3 \end{pmatrix}$$

$$A_7 = \begin{pmatrix} 2 & 0 & -2 \\ -3 & -1 & 2 \\ 2 & 0 & -2 \end{pmatrix}$$

$$A_8 = \begin{pmatrix} 2 & 4 & 3 \\ -2 & 2 & 0 \\ 3 & 0 & 2 \end{pmatrix}$$

$$A_9 = \begin{pmatrix} -1 & 2 & -2 \\ 2 & -1 & 2 \\ 2 & -2 & 3 \end{pmatrix}$$

31.- A matrix $A \in M_2$ verifies the following conditions: $A \begin{pmatrix} 1 \\ -1 \end{pmatrix} = \begin{pmatrix} 3 \\ 1 \end{pmatrix}$ and $(2, -1)$ is an eigenvector of A associated to the eigenvalue $\lambda = -2$. Find the matrix A indicating if it is diagonalizable or not. If affirmative, give a diagonal matrix D and an invertible matrix P such that $D = P^{-1}AP$.

32.- Let $A = \begin{pmatrix} 1 & a \\ 2 & b \end{pmatrix}$ with $a, b \in \mathbb{R}$. Calculate the values of the parameters a and b so that the vector $(2, -1)$ is eigenvector of A associated to the eigenvalue 2.

33.- Find a matrix $A \in M_3$ of eigenvalues $-1, 1, 2$ with associated eigenvectors $(1, 0, -1)$, $(1, 1, 0)$, $(3, -3, 1)$, respectively.

34.- Find a matrix $A \in M_3$ with eigenvalue $\lambda_1 = 1$ (simple) and eigenvalue $\lambda_2 = 2$ (double) with associated eigenvectors $(1,1,0)$, $(1,0,0)$, respectively.

35.- Find a matrix $A \in M_3$ with eigenvalue $\lambda_1 = -1$ (double) and eigenvalue $\lambda_2 = 3$ (simple) with associated eigenvectors $(1,0,2)$, $(-1,0,0)$ and $(0,1,1)$, respectively.

36.- Let $A = \begin{pmatrix} 3 & 2 \\ a & b \end{pmatrix}$ with $a, b \in \mathbb{R}$. Calculate the values of the parameters a and b so that A has eigenvalues 1 and -1. Is A a diagonalizable matrix?

37.- Consider the matrix $A = \begin{pmatrix} 1 & 0 & 3 \\ 3 & -2 & a \\ 3 & 0 & 1 \end{pmatrix}$ with $a \in \mathbb{R}$.

- a) For which values of the parameter a is $\lambda = -2$ an eigenvalue of A ?
- b) For which values of the parameter a is the matrix A diagonalizable?

38.- Determine a matrix $A \in M_3$ such that $A \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} -2 \\ 3 \\ 1 \end{pmatrix}$ and that his eigenvectors be

the non-null vectors of \mathbb{R}^3 belonging to the sets $\{(x, y, z) \in \mathbb{R}^3 \mid x = z\}$, $\{(x, y, z) \in \mathbb{R}^3 \mid x + y = 0, z = 0\}$.

39.- The matrix $A = \begin{pmatrix} a & 1 & p \\ b & 2 & q \\ c & -1 & r \end{pmatrix}$ admits as eigenvectors $(-1, -1, 0)$, $(1, 0, -2)$ and $(0, -1, 1)$

associated to the eigenvalues 3, 0 and 3/2 respectively. Answer the following items:

- a) Find the unknown elements of A .
- b) Is A diagonalizable? If affirmative, perform the diagonalization.

40.- Consider the matrix $A = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -1 & 0 & 1 & -1 \\ 0 & 1 & 2 & 0 \\ 0 & -1 & 1 & 1 \end{pmatrix}$

- a) Find its eigenvalues and eigenvectors. It is A diagonalizable?
- b) Check that that the determinant of the matrix A is the product of its eigenvalues.

41.- Check that the matrices $A = \begin{pmatrix} 2 & 2 & 1 \\ 1 & 3 & 1 \\ 1 & 2 & 2 \end{pmatrix}$ and $B = \begin{pmatrix} 2 & 1 & -1 \\ 0 & 2 & -1 \\ -3 & -2 & 3 \end{pmatrix}$ have the same

eigenvalues but, however, are not similar.

42.- Calculate A^{100} and, in general, A^k , with $k \in \mathbb{N}$, for the matrix $A = \begin{pmatrix} 0 & 2 \\ 1 & -1 \end{pmatrix}$.

43.- Calculate A^k with $k \in \mathbb{N}$ odd, for the matrix $A = \begin{pmatrix} 2 & -1 \\ 3 & -2 \end{pmatrix}$.

44.- Calculate $A^n \quad \forall n \in \mathbb{N}$ in each one of the following cases:

$$\text{a) } A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 2 \end{pmatrix} \quad \text{b) } A = \begin{pmatrix} 2 & -1 & 3 \\ 0 & 1 & -1 \\ 0 & 0 & 3 \end{pmatrix}$$

45.- Calculate a symmetric matrix $A \in M_3$ that verify: $v = (1, -1, 0)$ is eigenvector of A ,

$$A \begin{pmatrix} 2 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 2 \\ 0 \\ 2 \end{pmatrix} \text{ and } |A| = 0. \text{ Calculate } A^{50}.$$

46.- Determine for which values of the parameters $b, c \in \mathbb{R}$ the following matrices are diagonalizable. If affirmative, find a diagonal matrix similar to the given one.

$$A_1 = \begin{pmatrix} 5 & 0 & 0 \\ 0 & -1 & b \\ 3 & 0 & c \end{pmatrix} \quad A_2 = \begin{pmatrix} b & c & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{pmatrix} \quad A_3 = \begin{pmatrix} 1 & -2 & 7 \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix}$$

47.- For each one of the following matrices $A_i, i=1,2,3$, find, if it is possible, an invertible matrix P and a diagonal matrix D so that $D = P^{-1}A_iP$.

$$A_1 = \begin{pmatrix} -2 & 0 & 0 \\ 0 & -2 & -1 \\ 0 & -1 & -2 \end{pmatrix} \quad A_2 = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} \quad A_3 = \begin{pmatrix} 2 & 0 & 1 \\ 0 & 3 & 0 \\ 1 & 0 & 2 \end{pmatrix}$$

48.- The characteristic polynomial of a matrix A of order 3 is $P(\lambda) = -\lambda^3 + 21\lambda - 20$.

With this information, can you justify whether A is invertible and/or diagonalizable?

49.- The eigenvalues of a certain diagonalizable matrix $A \in M(n \times n)$ are the roots of the characteristic polynomial $P(\lambda) = \lambda^5 + \lambda^4 - 5\lambda^3 - 5\lambda^2 + 4\lambda + 4$.

- Determine the eigenvalues and their multiplicity.
- Determine the dimension of the matrix A and $\text{Rk}(A)$.
- Calculate, if possible, $|A^{-1}|$ and $|\frac{1}{2}A^{-1}|$.
- Calculate the eigenvalues of A^2 and their multiplicity.

50.- Consider the matrix $A = \begin{pmatrix} 1 & 1 & 0 \\ 2 & 0 & 0 \\ 1 & 1 & 2 \end{pmatrix}$

- Calculate the eigenvalues and eigenvectors of A .
- Is it diagonalizable? If it is, calculate a similar diagonal matrix D and a matrix P such that $D = P^{-1}AP$.
- Is there any value of the parameter a so that $(3, -6, a)$ is an eigenvector of A ?

51.- Let A be a matrix of order 3 whose characteristic polynomial is

$$P(\lambda) = -\lambda^3 - 2\lambda^2 + 5\lambda + 6 :$$

- Calculate the eigenvalues of A and the determinant of A .
- Explain if the following statements are true or false:
 - The homogeneous linear system, $AX = 0$, is consistent and determined.
 - There is a diagonal D matrix and a regular P matrix such that $D = P^{-1}AP$.

52.- Given a symmetric matrix A of order 4 whose eigenvalues are 1, 3, -5 and -5:

- Reason if A^{-1} exists and, if so, calculate its determinant.
- Obtain the rank of the matrix $A + 5I_4$?

53.- Let $A = \begin{pmatrix} m & 0 & 0 \\ 0 & 1 & -1 \\ 0 & -1 & 1 \end{pmatrix}$ with m being a real number:

- a) Study the existence of the inverse matrix of A according to the values of the parameter m. Calculate, when possible, A^{-1} .
- b) Determine the values of the parameter m so that A is diagonalizable.
- c) Calculate the characteristic polynomial and the eigenvalues of A.
- d) For $m = 0$, calculate a diagonal matrix D and a matrix P such that $D = P^{-1}AP$.

54.- Given the matrix $A = \begin{pmatrix} -1 & 0 & -3 \\ 3 & 3 & 3 \\ -3 & 0 & -1 \end{pmatrix}$,

- a) Calculate the eigenvalues of A.
- b) Calculate the eigenvectors of A.
- c) Is it diagonalizable? If it is, calculate a regular matrix P and a diagonal matrix D such that $D = P^{-1}AP$.
- d) Calculate the eigenvalues of A^3 .

QUADRATIC FORMS

1.- Consider the quadratic form $Q(\mathbf{x})$ of associated matrix (respectively)

$$A_1 = \begin{pmatrix} -1 & 1 & 0 \\ 1 & -2 & 1 \\ 0 & 1 & -1 \end{pmatrix} \quad A_2 = \begin{pmatrix} 3 & 2 & 4 \\ 2 & 0 & 2 \\ 4 & 2 & 3 \end{pmatrix} \quad A_3 = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} \quad A_4 = \begin{pmatrix} -2 & 0 & 0 \\ 0 & -2 & 1 \\ 0 & 1 & -2 \end{pmatrix}$$

Answer to the following points:

- Express $Q(\mathbf{x})$ in polynomial form.
- Find, by the method of eigenvalues, a diagonal expression for $Q(\mathbf{x})$.
- Classify $Q(\mathbf{x})$.

2.- Consider the quadratic form $Q(\mathbf{x})$ of associated matrix (respectively)

$$A_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 2 \\ 0 & 2 & 2 \end{pmatrix} \quad A_2 = \begin{pmatrix} 0 & -1 & 1 \\ -1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix} \quad A_3 = \begin{pmatrix} 3 & 2 & 4 \\ 2 & 0 & 2 \\ 4 & 2 & 3 \end{pmatrix}$$

Answer to the following points:

- Express $Q(\mathbf{x})$ in polynomial form.
- Find, by completing squares, a diagonal expression for $Q(\mathbf{x})$.
- Classify $Q(\mathbf{x})$.

3.- Consider the quadratic form $Q(x, y, z) = 2x^2 + 5y^2 + 5z^2 + 4xy - 4xz - 8yz$ and answer:

- Find a diagonal expression for Q .
- Classify Q .

4.- For each one of the following matrices, answer:

$$A_1 = \begin{pmatrix} 2 & 1 & 2 \\ 1 & 1 & 1 \\ 2 & 1 & 4 \end{pmatrix} \quad A_2 = \begin{pmatrix} 4 & 1 & 1 \\ 1 & 4 & 1 \\ 1 & 1 & 4 \end{pmatrix} \quad A_3 = \begin{pmatrix} -2 & -2 & 1 \\ -2 & 1 & -2 \\ 1 & -2 & -2 \end{pmatrix}$$

$$A_4 = \begin{pmatrix} 5 & -2 & 0 & -1 \\ -2 & 2 & 0 & 1 \\ 0 & 0 & 1 & -2 \\ -1 & 1 & -2 & 2 \end{pmatrix} \quad A_5 = \begin{pmatrix} -1 & 1 & 0 & 1 \\ 1 & -1 & 0 & 1 \\ 0 & 0 & -2 & 0 \\ 1 & 1 & 0 & -1 \end{pmatrix}$$

$$A_6 = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 4 & 1 \\ 1 & 1 & 4 \end{pmatrix} \quad A_7 = \begin{pmatrix} 2 & 2 & -2 \\ 2 & 3 & -4 \\ -2 & -4 & 0 \end{pmatrix}$$

5.- Classify by their sign the following quadratic forms:

- a) $Q(x, y) = 3x^2 - 4xy + 7y^2$.
- b) $Q(x, y) = x^2 + y^2 + 2xy$.
- c) $Q(x, y) = 6xy - 2x^2 - 5y^2$.
- d) $Q(x, y) = 4y^2 + 8xy$.
- e) $Q(x, y, z) = y^2 + 2xy + 2yz$.
- f) $Q(x, y, z) = x^2 + 4y^2 + z^2 + 2xy + 2xz + 4yz$
- g) $Q(x, y, z) = x^2 + 2y^2 + z^2 + xz + 2xy + 2yz$.
- h) $Q(x, y, z) = 4x^2 + 4y^2 + z^2 - 4xy$.
- i) $Q(x, y, z) = 2xy + 2xz + 2yz$.
- j) $Q(x, y, z) = x^2 + y^2 + z^2$.
- k) $Q(x, y, z) = -4x^2 + y^2 + 3z^2 + 3xz + yz$.
- l) $Q(x, y, z) = 2x^2 + 2xz + 3y^2 + 2z^2$.
- m) $Q(x, y, z) = 2x^2 - y^2 + 3z^2 - 3xy + 4yz - 2xz$.
- n) $Q(x, y, z) = x^2 + 10y^2 + 6xy$.
- o) $Q(x, y, z) = x^2 + 4y^2 + 3z^2 + 4xy + 2xz + 4yz$.
- p) $Q(x, y, z, t) = 2xz - 3yt + 2xt - t^2$.
- q) $Q(x, y, z) = x^2 - y^2 - 2z^2 + 2xy + 4yz$.
- r) $Q(x, y, z) = 2xy + 4yz - 4xz - x^2 - y^2 + 4z^2$.

6.- Show that for all $\forall x, y, z \in \mathbb{R}$ it holds $x^2 + y^2 + z^2 \geq xy + xz + yz$.

7.- Determine for what value of $\alpha \in \mathbb{R}$ the following quadratic forms are semidefinite indicating if it is positive or negative.

- a) $Q(x, y, z) = x^2 + 2y^2 + \alpha z^2 - 2xz$.
- b) $Q(x, y, z) = x^2 + \alpha y^2 + \alpha z^2 + 2yz$.

8.- Classify, depending on the values of $\beta \in \mathbb{R}$, the quadratic form of expression $Q(x, y, z, t) = x^2 + y^2 + z^2 + t^2 + 2\beta yt + 2\beta xz$.

9.- Study, depending on the values of the real parameter a , the sign of the quadratic form of associated matrix $A = \begin{pmatrix} a & 2 & 0 \\ 2 & a & 3 \\ 0 & 3 & 3 \end{pmatrix}$.

10.- Consider the quadratic form of associated matrix $A = \begin{pmatrix} -1 & a & 2 & 1 \\ a & 0 & 1 & 0 \\ 2 & 1 & 4 & 2 \\ 1 & 0 & 2 & 1 \end{pmatrix}$. Is it negative

definite for any value of the real parameter a ?

11.- Classify the following restricted quadratic forms:

a) $Q(x, y) = 2x^2 + y^2 + 2\sqrt{2}xy$ to $S = \{(x, y) \in \mathbb{R}^2 \mid x - \sqrt{2}y = 0\}$.

b) $Q(x, y, z) = x^2 + 4y^2 + 5z^2 + 2xy - 2xz + 4yz$ to the vectors (x, y, z) such that $x + 2y - z = 0$ and $2x - 3y + z = 0$.

c) $Q(x, y, z) = 2x^2 + y^2 - 4xy + 2yz$ to the vectors (x, y, z) such that $x - y + z = 0$.

d) $Q(x, y, z, t) = x^2 - z^2 + 2xz + xt + 2yz$ to the vectors (x, y, z, t) such that $x + y - z = 0, y - t = 0$.

12.- Classify the quadratic form $Q(x, y, z) = x^2 + y^2 - 2z^2$ restricted to:

a) $S = \{(x, y, z) \in \mathbb{R}^3 \mid x + z = 0\}$.

b) $S = \{(x, y, z) \in \mathbb{R}^3 \mid x = y = -z\}$.

13.- Classify $Q(x, y, z) = 2x^2 + 2y^2 + 2z^2 + 2xy + 2xz + 2yz$ restricted to:

a) $S = \{(x, y, z) \in \mathbb{R}^3 \mid x - 2z = 0\}$.

b) $S = \{(0, 0, z) \mid z \in \mathbb{R}\}$.

14.- Classify the quadratic form $Q(x, y, z) = x^2 + y^2 + z^2 + 2xy + 2xz + 2yz$ restricted to the set $S = \{(x, y, z) \in \mathbb{R}^3 \mid x - y - 2z = 0\}$.

15.- Consider the quadratic form $Q(x, y, z) = 4x^2 + 5y^2 + z^2 - 4xz$.

- a) Find a diagonal expression for Q .
- b) Classify the sign of Q .
- c) Classify Q restricted to $S = \{(x, y, z) \in \mathbb{R}^3 \mid x - z = 0\}$.

16.- Consider the quadratic form of associated matrix $A = \begin{pmatrix} 1 & 0 & b \\ 0 & 2 & 0 \\ b & 0 & 1 \end{pmatrix}$, where b is a real

parameter.

- a) Determine its sign as a function of the values of the parameter b .
- b) Determine its sign restricted to the vectors (x, y, z) satisfying $y = 2z$, for any value of b .

17.- Let $Q(x, y, z) = y^2 - xy - xz - yz$.

- a) Find a diagonal expression for Q .
- b) Classify Q .
- c) Classify, depending on the values of the real parameter α , Q restricted to the set $S = \{(x, y, z) \in \mathbb{R}^3 \mid x = y = \alpha z\}$.

18.- Let $Q(x, y, z) = 2ax^2 + y^2 + z^2 + 4axz$, with $a \in \mathbb{R}$.

- a) Classify Q depending on the values of the real parameter a .
- b) If $a = -1$, find subsets S_1 and S_2 of \mathbb{R}^3 such that Q restricted to S_1 be positive definite and Q restricted to S_2 be negative definite.

19.- Considering the quadratic form $Q(x, y, z)$ represented by the matrix $A = \begin{pmatrix} -2 & 3 & 0 \\ 3 & -2 & 0 \\ 0 & 0 & 5 \end{pmatrix}$

answer reasonably the following questions:

- a) Does the quadratic form Q admit the expression $Q(\tilde{x}, \tilde{y}, \tilde{z}) = -2\tilde{x}^2 - 2\tilde{y}^2 - 5\tilde{z}^2$?
- b) Justify if there is any $(x_0, y_0, z_0) \in \mathbb{R}^3$ for which it is verified that $Q(x_0, y_0, z_0) > 0$.
- c) Define, if possible, a subset \mathbb{R}^3 where the quadratic form Q is definite negative.

20.- Given the quadratic form $Q(x, y, z) = 2x^2 - y^2 - 4z^2 + 4zy$. It is requested:

- a) Calculate a diagonal expression for Q .
- b) Justify if there exist some $(x_0, y_0, z_0) \in \mathbb{R}^3$ such that $Q(x_0, y_0, z_0) < 0$.
- c) Study the sign of Q constrained to $S = \{(x, y, z) \in \mathbb{R}^3 \mid y - 2z = 0\}$.

21.- Given the quadratic form $Q(x, y, z) = x^2 + 3y^2 + z^2 + 2xz$:

- a) Classify $Q(x, y, z)$.
- b) Can $Q(\bar{x}, \bar{y}, \bar{z}) = \bar{x}^2 + 3\bar{y}^2 + 2\bar{z}^2$ be a diagonal expression of $Q(x, y, z)$?

22.- Given the high value of the public deficit of an imaginary country, the government decides to create a new tax T whose amount is a function of the payments (or reimbursements if applicable) of the income tax, R , and of the payments of the heritage tax, P , in such way that $T = 2R^2 + 4P^2 - 4RP$. The government, before setting up the new tax, wants to make sure that no taxpayer will be reimbursed by this new tax. Check that the new tax will fulfil its purpose with all and each one of the taxpayers.

23.- The economists of a company claim that the production function is of the type: $P = L^2 + K^2 - 2LK$, being L and K the number of workers and of machines, respectively. Besides that, it is known that each machine needs two workers for it to work. Check that indeed, P is a production function with the condition given.

24.- An investor estimates that by investing in three securities, A, B and C, the resulting portfolio will have a profitability of $U(x, y, z) = 2x^2 - 2y^2 - 7z^2 + 2xy + 6xz$, being x , y and z , the returns at the time of purchase of each value A, B and C respectively.

- a) Analyze if said portfolio can generate losses.
- b) Analyze how the previous response affects an economic situation in which the three products have the same profitability.
- c) If the investor knows that the profitability of the product A is double of that of B, Will there be gains?

FUNCTIONS FROM \mathbb{R}^n TO \mathbb{R}^m

Note: it will be understood that $\log x = \log_{10} x$ and $\ln x = \log_e x$

1.- Determine and represent graphically the domain of the following functions:

a) $f(x) = \sqrt{x^2 - 16}$

b) $f(x) = \frac{1}{x^2 - 1}$

c) $f(x) = \frac{1}{\sqrt{x-3}}$

d) $f(x) = \frac{1}{\sqrt{1-x^2}}$

e) $f(x) = \frac{\sqrt[3]{x+1}}{x^2 + x + 1}$

f) $f(x) = e^{\sqrt{\frac{x+1}{x-3}}}$

g) $f(x) = \sqrt{-x} + \frac{1}{\sqrt{2+x}}$

h) $f(x) = \ln(x^2 - 4)$

i) $f(x) = \ln \frac{x^2 - 3x + 2}{x + 1}$

j) $f(x) = \ln(x + 2) + \ln(x - 2)$

k) $f(x, y) = 3x + y$

l) $f(x, y) = \frac{3x + y}{x^2 + 2y^2}$

m) $f(x, y) = \frac{1}{\sqrt{x-y}}$

n) $f(x, y) = \frac{y}{x^2}$

o) $f(x, y) = \ln(2x - y + 1)$

p) $f(x, y) = x + \sqrt{y}$

q) $f(x, y) = \sqrt{1-x^2} + \sqrt{y^2-1}$

r) $f(x, y) = \sqrt{1-x^2-y^2}$

s) $f(x, y) = \frac{1}{\sqrt{x^2 + y^2 - 1}}$

t) $f(x, y) = \ln(3x^2 + y^2 + 4)$

u) $f(x, y, z) = \sqrt{1-x^2-y^2-z^2}$

v) $f(x, y) = \left(\ln(x+y), \sqrt{4-x^2-y^2} \right)$

2.- Represent graphically the level (contour) curves of the following functions:

a) $f(x, y) = x + y$

b) $f(x, y) = (x + y)^2$

c) $f(x, y) = x^2 + y^2$

d) $f(x, y) = \frac{x}{y}$

e) $f(x, y) = \sqrt{xy}$

f) $f(x, y) = x^y, x > 0$

g) $f(x, y) = \frac{y}{x^2}$

h) $f(x, y) = \ln(2x - y + 1)$

i) $f(x, y) = \frac{x+y}{x}$

3.- Compute the following limits if possible:

- a) $\lim_{x \rightarrow 1} \frac{x^2 - 1}{2x^2 - x - 1}$ b) $\lim_{x \rightarrow 1} \frac{x - 1}{\sqrt{x^3 + 3} - 2}$ c) $\lim_{x \rightarrow 0} \left(\frac{1+x}{1-x} \right)^x$
- d) $\lim_{x \rightarrow 0} \frac{x^2}{\frac{1}{1+2^x}}$ e) $\lim_{x \rightarrow 0} \frac{2}{\frac{1}{3+4^x}}$ f) $\lim_{x \rightarrow +\infty} \frac{5x^2 - 6x}{3x^2 - 8}$
- g) $\lim_{x \rightarrow +\infty} \left(1 - \frac{2}{3x} \right)^x$ h) $\lim_{x \rightarrow +\infty} \left(\frac{2x+1}{2x+4} \right)^{\frac{x^2}{x+1}}$ i) $\lim_{x \rightarrow +\infty} \frac{x^5}{e^{-x}}$
- j) $\lim_{x \rightarrow +\infty} \left(\frac{2x+1}{x} \right)^x$ k) $\lim_{x \rightarrow +\infty} \left(\frac{5x+1}{8x-1} \right)^{-x}$ l) $\lim_{(x,y) \rightarrow (1,2)} \frac{3x^2y}{4x - y + 1}$
- m) $\lim_{(x,y) \rightarrow (0,0)} \frac{3x^2}{5x^2 + y^2}$ n) $\lim_{(x,y) \rightarrow (0,1)} \left(x^2 + 5xy - 8, e^{x^2y}, \ln(2x + y) \right)$

4.- Study the continuity of the following functions:

- a) $f(x) = \begin{cases} \frac{e^x}{x-1} & \text{if } x \leq 0 \\ x^3 + 1 & \text{if } x > 0 \end{cases}$ b) $f(x, y) = \begin{cases} \frac{2x^2 - 3y^2}{7x^2 + y^2} & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0) \end{cases}$

5.- Compute the first derivative of the following functions:

- a) $f(x) = e^5$ b) $f(x) = \frac{e^x}{x}$
- c) $f(x) = \frac{1}{x}$ d) $f(x) = e^{\frac{-x}{2}}$
- e) $f(x) = \frac{1+x}{1-x}$ f) $f(x) = \sqrt{2x-1}$
- g) $f(x) = \ln x + \sqrt{x^2 + 1}$ h) $f(x) = \frac{\pi}{x} + \ln 2$
- i) $f(x) = \frac{2}{2x-1} - \frac{1}{x}$ j) $f(x) = \frac{1+\sqrt{x}}{1-\sqrt{x}}$
- k) $f(x) = \ln x \log x - \ln a \log_a x$ l) $f(x) = \frac{3}{56(2x-1)^7} - \frac{1}{24(2x-1)^6} - \frac{1}{40(2x-1)^5}$
- m) $f(x) = \sqrt[3]{2e^x - 2^x} + 1 + \ln^5 x$ n) $f(x) = x^2 10^{2x}$

o) $f(x) = \ln^2 x - \ln(\ln x)$

p) $f(x) = \frac{x}{2}\sqrt{x^2 - a^2} - \frac{a^2}{2}\ln(x + \sqrt{x^2 - a^2})$

q) $f(x) = x^4(a - 2x^3)^2$

r) $f(x) = \sqrt{(x+a)(x+b)(x+c)}$

s) $f(x) = \sqrt[3]{x + \sqrt{x}}$

t) $f(x) = (2x+1)(3x+2)\sqrt[3]{3x+2}$

u) $f(x) = \frac{4}{3}\sqrt[4]{\frac{x-1}{x+2}}$

v) $f(x) = \ln(\sin x)$

w) $f(x) = \sin x^4$

x) $f(x) = \cos^4 x$

y) $f(x) = \sin^2 x \cos x^6$

6.- Compute the partial derivatives of the following functions:

a) $f(x, y, z) = e^{\frac{x}{y}} + e^{\frac{z}{y}}$

b) $f(x, y) = \ln \frac{\sqrt{x^2 + y^2} - x}{\sqrt{x^2 + y^2} + x}$

c) $f(x, y, z) = e^{x^2 + y^2 + z^2}$

d) $f(x, y) = \sqrt{xy + \frac{x}{y}}$

e) $f(x, y) = x^3 + y^3 - 3axy$

f) $f(x, y) = \frac{x-y}{x+y}$

g) $f(x, y) = \ln(x + \sqrt{x^2 + y^2})$

h) $f(x, y) = x^y$

i) $f(x, y, z) = z^{xy}$

j) $f(x, y) = \frac{x+y}{\sqrt[3]{x^2 + y^2}}$

k) $f(x, y) = x^2 \sin^2 y$

l) $f(x, y) = \frac{e^{ax}(\sin x + a \cos y)}{a^2 + b^2}$

7.- Check that $y = xe^{-x}$ verifies the equation $x \frac{dy}{dx} = (1-x)y$.

8.- Compute y' starting from the following expressions:

a) $2x^2 + 5xy + y^2 = 19$

b) $x^2 + y^2 = 25$

c) $y = 1 + xe^y$

d) $\ln y + e^{\frac{y}{x}} = 8$

e) $\ln y + \frac{x}{y} = 7$

f) $x^y = y^x$

9.- Find the equation of the tangent straight line to the graph of the function $f(x) = \frac{1}{x}$ at $x=1$. Find an approximate value of $f(1.1)$.

10.- Compute the following limits:

a) $\lim_{x \rightarrow 0} \frac{\sin x}{x}$

b) $\lim_{x \rightarrow 0} \frac{2(1 - \cos x)}{x^2}$

c) $\lim_{x \rightarrow 0} \frac{e^x - 1}{x}$

d) $\lim_{x \rightarrow 0} \frac{\sin 3x^2}{3x^2}$

e) $\lim_{x \rightarrow 0} \frac{1 - \cos \sqrt{x}}{\frac{1}{2}x}$

f) $\lim_{x \rightarrow 0} \frac{1 - \cos(\sin x)}{x^2}$

g) $\lim_{x \rightarrow 0} \frac{\ln(1 + x^2)}{x^2}$

h) $\lim_{x \rightarrow 0} \frac{e^{x^2} - 1}{x^2}$

i) $\lim_{x \rightarrow 0} \frac{\ln(1 + \sin x)}{\sin x}$

j) $\lim_{x \rightarrow 0} \frac{e^{2x} - 1}{x^3}$

k) $\lim_{x \rightarrow 0} \frac{\sin x \operatorname{tg} x}{1 - \cos x}$

l) $\lim_{x \rightarrow 0} \frac{1 - \cos 4x}{5x^2}$

m) $\lim_{x \rightarrow 0} \frac{\ln(1 + x)}{1 - e^x}$

n) $\lim_{x \rightarrow 0} \frac{\sin^2 x}{1 - \cos 2x}$

o) $\lim_{x \rightarrow 0} \frac{4(x - \ln(1 + x))}{x \ln(1 + x)}$

p) $\lim_{x \rightarrow 0^+} x^{\frac{1}{x}}$

q) $\lim_{x \rightarrow 0^+} (\cos x)^{\frac{1}{x}}$

r) $\lim_{x \rightarrow +\infty} x [\ln(x + 1) - \ln x]$

s) $\lim_{x \rightarrow +\infty} \frac{1}{x} \ln x$

t) $\lim_{x \rightarrow +\infty} \frac{\ln x}{x^2}$

u) $\lim_{x \rightarrow +\infty} \frac{\ln(1 + x^6)}{x^2}$

v) $\lim_{x \rightarrow +\infty} (\ln x^3) e^{-x}$

11.- Compute $\nabla f(x, y)$ when $f(x, y) = x^2 y + y^3$.

12.- Find the Jacobian matrix of the following functions:

a) $f(x) = \ln^2 x$

b) $f(x_1, x_2) = (x_1 - x_2)^2$

c) $f(x_1, x_2) = (x_1 + x_2, x_1 - x_2)$

d) $f(t) = (t, e^t, e^{-t})$

13.- Compute $\frac{dz}{dt}$ in the functions:

a) $z = 3x + y$, with $x = t^2 + 1$; $y = e^t$.

b) $z = xy + yu + xu$, with $x = t$; $y = e^{-t}$; $u = \ln(t)$.

c) $z = e^{xy}$, with $x = t \cos t$; $y = t \sin t$.

14.- Compute $\frac{dz}{du}(1)$ in the function $z = 3x^2 + 2xy - y^2$, with $x = u^2 + 3u$; $y = 2u^2 - u$.

15.- Compute $\frac{\partial u}{\partial r}$ and $\frac{\partial u}{\partial s}$ in the function $u = z \sin \frac{y}{x}$, with $x = 3r^2 + 2s$; $y = 4r - 2s^3$, $z = 2r^2 - 3s^2$.

16.- Compute $\frac{\partial z}{\partial x}$, $\frac{\partial z}{\partial y}$ and evaluate them at the point $(x, y) = (0, 1)$, in the following cases:

a) $z = u + v$, with $u = x + e^y$; $v = \ln(y) + e^{-x}$.

b) $z = \frac{\sin u}{v}$, with $u = s - t$; $v = s + x$; $s = y^2 - x$; $t = e^y$.

17.- Prove that $x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = 2$, if $z = \ln(x^2 + xy + y^2)$.

18.- Prove that $x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = xy + z$, if $z = xy + x e^{y/x}$.

19.- Prove that it holds

$$(x^2 - y^2) \frac{\partial z}{\partial x}(x, y) + xy \frac{\partial z}{\partial y}(x, y) = xyz,$$

where $z = e^y F\left(y e^{\frac{x^2}{2y^2}}\right)$ and F being a differentiable real function of a real variable.

20.- Let $f(x, y) = g(x^2 + y^2)$, being $g : \mathbb{R} \rightarrow \mathbb{R}$ a differentiable function at any point of the real line. If $g'(5) = 3$, compute $\nabla f(1, 2)$.

21.- Determine the differential of the following functions at the indicated points:

a) $y = \ln x$, at 2

b) $y = \frac{x^2}{e^x}$, at 0

c) $z = \sqrt{x^2 + y^2}$, at $(1, 1)$

d) $u = \left(xy + \frac{x}{y}\right)^z$, at $(1, 1, 1)$.

e) $z = \ln \frac{2x}{y^2}$, at $(1, e)$

j) $f(x, y) = x^2 e^{\frac{y}{x}}$, at (x, y) with $x \neq 0$

k) $z = (e^{x+y} + y, xy^2)$, at $(0, 0)$

22.- Let $f(x) = \sqrt[3]{x}$, using the concept of differential, study the approximate variation of the function when we change $x=27$ into $x=26.9$.

23.- Compute the directional derivative of the following functions in the indicated direction and at the indicated point:

a) $f(x,y) = 3 - 2x^2 + y^3$, in the direction of $v = \left(\frac{1}{2}, \frac{-\sqrt{3}}{2}\right)$ at the point $P=(1,2)$

b) $f(x,y) = x^2 + y + 1$, in the direction $\left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$ at the point $(0,0)$

c) $f(x,y) = \log(x^2 + y^3)$, in the direction of $v = \left(\frac{2}{\sqrt{13}}, \frac{3}{\sqrt{13}}\right)$ at the point $P = (1,3)$

d) $f(x,y,z) = x \sin(yz)$, in the direction $\left(\frac{1}{\sqrt{6}}, \frac{2}{\sqrt{6}}, -\frac{1}{\sqrt{6}}\right)$ at the point $(1,3,0)$

e) $f(x,y,z) = x \sin y + y \cos z + z \sin x$, in the direction $\left(\frac{\sqrt{2}}{2}, 0, \frac{\sqrt{2}}{2}\right)$ at the point $(0,0,0)$

f) $f(x,y,z) = (x^2 + yz^2, \sin(x^2 + y^2))$, in the direction $\left(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}\right)$ at the point $(\sqrt{\pi}, \sqrt{\pi}, 1)$

24.- Let $f(x,y)$ be a differentiable function at the point $(1,2)$. Knowing that $f_v(1,2) = 5$ and $f_w(1,2) = 6$ with $v = \left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$ and $w = \left(\frac{1}{\sqrt{5}}, -\frac{2}{\sqrt{5}}\right)$, compute $\nabla f(1,2)$.

25.- Given the vector field $f(x,y) = (xy, x^2, y^2)$,

a) Study its differentiability at the point $(-1,2)$.

b) In the case of being differentiable, compute its differential at that point.

c) Find f_v' at the point $(-1,2)$ being $v = \left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$.

26.- Compute y'' in $y = \ln(x + \sqrt{4 + x^2})$.

27.- Given $f(x,y) = x e^{2y-x}$, compute $\frac{\partial f}{\partial x}$, $\frac{\partial f}{\partial y}$, $\frac{\partial^2 f}{\partial x^2}$, $\frac{\partial^2 f}{\partial y^2}$, $\frac{\partial^2 f}{\partial x \partial y}$ and $\frac{\partial^2 f}{\partial y \partial x}$.

28.- Compute the Hessian matrix of the following functions:

- a) $f(x, y) = x^4 + y^4 + 4xy - 2x^2 - 2y^2$.
- b) $f(x, y) = x^2 + y^2 + xy - 4\ln x - 10\ln y, x, y > 0$.
- c) $f(x, y) = 9x^2 + y^2 + 6xy + 12x + 4y$.
- d) $f(x, y) = x^3 + y^3 - 9xy + 27$.
- e) $f(x, y, z) = x^2 + y^2 + z^2 - xy + x - 2z$.
- f) $f(x, y, z) = xyz$.

29.- If F and G are real functions of a real variable with continuous second order derivatives, prove that the function $z = x F\left(\frac{y}{x}\right) + G\left(\frac{y}{x}\right)$ satisfies the following equation:

$$x^2 \frac{\partial^2 z}{\partial x^2}(x, y) + 2xy \frac{\partial^2 z}{\partial x \partial y}(x, y) + y^2 \frac{\partial^2 z}{\partial y^2}(x, y) = 0.$$

30.- Knowing that f and g are real functions of a real variable with derivatives of the second order and $z = f(x^2 + y^2) + g(x^2 + y^2)$, compute:

$$\frac{1}{x^2} \frac{\partial^2 z}{\partial x^2}(x, y) - \frac{1}{y^2} \frac{\partial^2 z}{\partial y^2}(x, y) - \frac{1}{x^3} \frac{\partial z}{\partial x}(x, y) + \frac{1}{y^3} \frac{\partial z}{\partial y}(x, y).$$

31.- Check that $z = \log(x^2 + y^2)$ satisfies the equation $\frac{\partial^2 z}{\partial x^2} + \frac{\partial^2 z}{\partial y^2} = 0$.

32.- If f is a real function of a real variable with second derivative, check that the function $z = x f(x + y) + y f(x + y)$ satisfies the equation:

$$\frac{\partial^2 z}{\partial x^2} - 2 \frac{\partial^2 z}{\partial x \partial y} + \frac{\partial^2 z}{\partial y^2} = 0.$$

33.- Let $f, g : \mathbb{R} \rightarrow \mathbb{R}$ be real functions of a real variable and $h(x, y) = f(y \cdot g(x))$. Assuming that there exist f'' and g'' in all of \mathbb{R} , compute $\frac{\partial^2 h}{\partial x^2}(x, y), \frac{\partial^2 h}{\partial y^2}(x, y)$.

34.- Compute the Taylor expansion of order 2 of the following functions in a neighbourhood of the indicated points:

- a) $f(x) = \ln(1 + x), x = 0$
- b) $f(x) = e^x, x = 0$

$$c) f(x) = \frac{1}{1+x}, \quad x = 0$$

$$d) f(x) = \frac{1}{(1+x)^2}, \quad x = 0$$

$$e) f(x) = \sqrt{1+x}, \quad x = 0$$

$$f) f(x) = \sin x, \quad x = 0$$

$$g) f(x) = \cos x, \quad x = 0$$

$$h) f(x) = \frac{2x^2 - 3}{(x-1)^2}, \quad x = 0$$

$$i) f(x) = \ln(x - 2x^2), \quad x = \frac{1}{3}$$

$$j) f(x) = \ln\left(\frac{1+x}{1-x}\right), \quad x = 0$$

$$k) f(x) = \ln\left(x + \sqrt{1+x^2}\right), \quad x = 1$$

$$l) f(x) = \sqrt[3]{6+x}, \quad x = 2$$

$$m) f(x) = (1+x)e^{-x}, \quad x = 0$$

$$n) f(x) = (1+e^x)^3, \quad x = \frac{1}{2}$$

$$o) f(x) = \sin^2 2x, \quad x = 0$$

$$p) f(x) = \ln(2x) - \frac{1}{x-1}, \quad x = 2$$

$$q) f(x) = e^x \ln(1-x), \quad x = 0$$

$$r) f(x) = \ln(9-x^2), \quad x = 0$$

$$s) f(x, y) = e^{2x-3y}, \quad (x, y) = (0, 0)$$

$$t) f(x, y) = \frac{x^2 y}{x^2 + y^2}, \quad (x, y) = (1, 1)$$

$$u) f(x, y, z) = x + yz + e^y, \quad (x, y, z) = (1, 0, 1)$$

35.- Given the function $f(x, y) = \sqrt{x+y} \ln y$, we consider the equation $f(x, y) = 2e$:

- Prove that this equation defines y as an implicit function of x in a neighbourhood of the point $(0, e^2)$.
- Compute $y'(0)$.

36.- Given the equation $e^z \sin(x+y) + e^y \sin(x+z) + e^x \sin(y+z) = 0$:

- Prove that it defines z as an implicit function of x and y in a neighbourhood of the point $(\pi, 0, \pi)$.
- Compute $\frac{\partial z}{\partial x}(\pi, 0)$ and $\frac{\partial z}{\partial y}(\pi, 0)$.

37.- Prove that the equation $x^2 y + xy^2 = 16$ defines y as an implicit function of x in a neighbourhood of the point $(2, 2)$. Is $y(x)$ increasing or decreasing at $x=2$?

38.- Consider the equation $3\alpha x^2 - \ln(yz) - \frac{3\alpha x}{yz} = 0$. For which values of the parameter α we can assure that the previous equation defines implicitly $x = x(y, z)$ in a neighbourhood of the point $(1, 1, 1)$?

39.- Given the equation $z \sin x - y \sin z = 0$:

a) Prove that it defines z as an implicit function of (x, y) in a neighbourhood of

$$\left(\frac{\pi}{2}, \frac{\pi}{2}, \frac{\pi}{2}\right).$$

b) Find the Taylor polynomial of degree 1 in a neighbourhood of the point $\left(\frac{\pi}{2}, \frac{\pi}{2}\right)$ of the function defined in a).

40.- Let $h : \mathbb{R}^2 \rightarrow \mathbb{R}$ be given by $h(x, y) = \sin(x^2 + y) + xy + a^2 y$.

a) For which values of a the equation $h(x, y) = 0$ defines y as an implicit function of x , $y = \varphi(x)$, in a neighbourhood of $(0, 0)$?

b) Compute $\varphi'(0)$ when possible.

c) Does it define the same equation x as an implicit function y in a neighbourhood of $(0, 0)$ for some value of a ?

d) Let $F(x, t) = (e^{x+t} + x^2 - 1, e^{\varphi(x)} + t \cos x - 1)$, with $\varphi(x)$ the implicit function of a). Prove that $JF(0, 0)$ is a regular matrix.

41.- Given the equation $e^{x^2 - y^2} + \alpha(x^2 + y^2) = 1 + \alpha$:

a) For which values of $\alpha \in \mathbb{R}$ does it define $y = y(x)$ as an implicit function in a neighbourhood of the point $\left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$?

b) For the previous values of $\alpha \in \mathbb{R}$, compute $\frac{dy}{dx}\left(\frac{1}{\sqrt{2}}\right)$.

42.- Let $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ be given by $f(x, y, z) = \alpha yz - y \ln(1 + z^2) + z \cos(x + 2y) + 1$, $\alpha \in \mathbb{R}$.

a) Prove that $f(x, y, z) = 0$ defines z as an implicit function of x and y in a neighbourhood of the point $(\pi, 0, 1)$ for any value of α .

b) Compute α such that $\frac{\partial z}{\partial x}(\pi, 0) = 0 = \frac{\partial z}{\partial y}(\pi, 0)$.

43.- Given the equation $xy^2 - yx^2 + z^2 \cos(xz) = 1$:

- Prove that it defines $z(x, y)$ as an implicit function in a neighbourhood of the point $(0, \sqrt{2}, 1)$.
- Find the tangent plane of $z(x, y)$ at the point $(0, \sqrt{2})$.
- Compute the directional derivative of $z(x, y)$ at $(0, \sqrt{2})$ respect to the direction $v = \left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}} \right)$.

44.- Let $F : \mathbb{R}^4 \rightarrow \mathbb{R}$ be given by $F(x, y, z, t) = x^3 z + y^3 t^2 - 1$:

- Prove that the equation $F(x, y, z, t) = 0$ defines t as an implicit function of the variables (x, y, z) in a neighbourhood of the point $(0, 1, 0, 1)$.
- If $t = \varphi(x, y, z)$ is the implicit function of a), compute $\nabla \varphi(0, 1, 0)$.

45.- Study whether the following functions are homogeneous, indicating the degree of homogeneity if in the affirmative:

a) $f(x, y) = \frac{\sqrt{x^4 + y^4}}{x}$

b) $f(x, y) = 3x^4 + 4x^2y^2 + 5y^4$

c) $f(x, y) = \frac{x^2y^2}{x^3 + y^3}$

d) $f(x, y) = \frac{2xy}{x^2 + y^2}$

e) $f(x, y) = (x^2 + y^2)^{-1/3}$

f) $f(x, y) = x^2 + 3xy^2 - 15x - 12y$

g) $f(x, y) = 90x^{1/3}y^{1/3}$

h) $f(x, y) = x^a y^b$

i) $f(x, y, z) = \left(\frac{x^3y + x^2yz - 4xz^3}{x - 2y} \right)^5$

j) $f(x, y, z) = \frac{1}{x^2} \ln \frac{y}{z}$

k) $f(x, y, z) = \sqrt[3]{x^2 - y^2} + \sqrt{x + z}$

l) $f(x, y, z) = \sqrt[5]{\frac{x^6 + y^4x^2 + yz^5}{2z^3}}$

m) $f(x, y, z) = \ln \frac{x - 2y}{y + 3z}$

n) $f(x, y, z) = e^{3x+y} + \sqrt[3]{xz}$

o) $f(x, y, z) = e^{\sqrt{\frac{x^2}{yz}}}$

p) $f(x, y, z) = x \ln \frac{y}{z} + \sqrt{2yz} + 7$

46.- For the functions of the previous exercise, compute the partial derivatives. Check that Euler's Theorem holds in the appropriate cases.

47.- Given $f(x, y) = x^4 y^2 e^{y/x}$, check that $x \frac{\partial f}{\partial x} + y \frac{\partial f}{\partial y} = 6f$. What can be deduced from the previous equality?

48.- Let f be an homogeneous function of degree m , such that $f(-1, 1) = 1$ and $f(-2, 2) = 1$. Compute m .

49.- Let f be a differentiable, homogeneous function of degree 2 with $\frac{\partial f}{\partial x}(3, 2) = 3$ and

$$\frac{\partial f}{\partial y}(3, 2) = 4. \text{ Compute } f(3, 2).$$

50.- Let $f(x, y, z)$ be a differentiable, homogeneous function of degree 3 such that the components of its gradient vector at $(1, 2, 3)$ are $(5, 2, 2)$. What is the value of the function at the point $(1, 2, 3)$?

51.- Let $z = z(x, y)$ be such that it verifies the equation $F\left(\frac{y}{x}, \frac{z}{x}\right) = 0$, F being a function with derivatives of first order, the second of them being not zero.

a) Prove that $z = z(x, y)$ is homogeneous of degree 1.

b) It is homogeneous $\frac{\partial z}{\partial x}$? If in the affirmative, of what degree? Reason out the answer.

52.- Let $f(x, y)$ be homogenous of degree 1 with derivativeds of second order. Prove that:

$$\frac{1}{y^2} \frac{\partial^2 f}{\partial x^2} = -\frac{1}{xy} \frac{\partial^2 f}{\partial x \partial y} = \frac{1}{x^2} \frac{\partial^2 f}{\partial y^2}.$$

53.- Let $f(x, y) = \frac{x^2 y^2}{x^2 + y^2}$. Prove that it holds:

$$x^2 \frac{\partial^2 f}{\partial x^2} + y^2 \frac{\partial^2 f}{\partial y^2} = 2 \frac{x^2 y^2}{x^2 + y^2} - 2xy \frac{\partial^2 f}{\partial x \partial y}.$$

54.- Given the function $f(x, y, z) = e^{\frac{\operatorname{tg} x^2 + y^2 + z^2}{xy}}$:

a) Check that it is homogeneous of degree 0.

b) Compute $x \frac{\partial f}{\partial x}(x, y, z) + y \frac{\partial f}{\partial y}(x, y, z) + z \frac{\partial f}{\partial z}(x, y, z)$.

55.- Let $f, g : \mathbb{R}^3 \rightarrow \mathbb{R}$ be differentiable functions and homogeneous of degrees 4 and 1, respectively. Prove that if $h(x, y, z) = f(x, y, z) \cdot g(x, y, z)$, it is verified:

$$x \frac{\partial h}{\partial x}(x, y, z) + y \frac{\partial h}{\partial y}(x, y, z) + z \frac{\partial h}{\partial z}(x, y, z) = 5 h(x, y, z).$$

56.- Let $f(x, y)$ be differentiable such that $\frac{\partial f}{\partial x}(1, 1) = 1$, $\frac{\partial f}{\partial y}(1, 1) = 1$, $f(1, 1) = 2$, $\frac{\partial f}{\partial x}(1, 2) = 1$,

$\frac{\partial f}{\partial y}(1, 2) = 1$, $f(1, 2) = 5$. Say whether f is homogeneous and if in the affirmative, of what degree.

57.- Let $f(x, y) = e^{x^2 + y^2}$.

- Compute the contour curves of $f(x, y)$ and represent them graphically.
- Compute $\nabla f(2, 1)$.
- Compute the directional derivative of f at the point $(2, 1)$ following the direction $\left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}} \right)$.
- Taking into account $\begin{cases} x = 2 + \ln t^2 \\ y = e^{t^3 - 1} \end{cases}$, compute $\frac{df}{dt}(1)$.

58.- Let the function $f(x, y) = \ln(1 + x^2 + 2y^2)$.

- Compute the domain of f .
- Compute the gradient vector of f at (x, y) .
- Is the function f differentiable?
- Compute the directional derivative of f respect to the direction of $\left(\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}} \right)$ at the point $(1, 1)$.
- Does the function f verify the conditions of Schwarz's Theorem?
- Using the result of e), compute the Hessian matrix of f at (x, y) .

59.- Let the function $f(x, y) = \ln\left(\frac{y}{x^2} + 1\right) - 1$.

- Determine the contour curves of the function and represent them graphically.
- Compute $\nabla f(1, 2e)$.
- Given the vector $v = \left(\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}\right)$, compute the directional derivative $f_v(1, 2e)$.
- Prove that the contour curve of level zero of f defines y as an implicit function of x . Taking the implicit differentiation, compute $y'(1)$.

60.- Consider the function of two variables $f(x, y) = \frac{y}{x + 2y}$.

- Determine its domain and represent it graphically.
- Determine and represent the contour curves of f .
- Prove that $\lim_{(x,y) \rightarrow (0,0)} f(x, y)$ does not exist.
- Determine the pairs $(a, b) \in \mathbb{R}^2$ such that the equation $f(x, y) = \frac{b}{a + 2b}$ defines y as an implicit function of x in a neighbourhood of the point (a, b) . For each of the previous pairs, compute $y'(a)$ by implicit differentiation.

61.- Let $F(x, y, z) = x^2 z y + e^{xz} - z^2 y + 4y$:

- Without computing the partial derivatives, reason out whether the following relation is true:

$$x \frac{\partial F}{\partial x}(x, y, z) + y \frac{\partial F}{\partial y}(x, y, z) + z \frac{\partial F}{\partial z}(x, y, z) = 4F(x, y, z).$$

- Compute $\nabla F(0, 2, 1)$.
- Compute the directional derivative of $F(x, y, z)$ with respect to the direction $\left(\frac{1}{\sqrt{3}}, \frac{-1}{\sqrt{3}}, \frac{-1}{\sqrt{3}}\right)$ at the point $(0, 2, 1)$.
- Prove that the equation $F(x, y, z) = 7$ defines x as an implicit function of y, z ($x = f(y, z)$) in a neighbourhood of the point $(0, 2, 1)$.
- Compute $\nabla f(2, 1)$.

- Compute the directional derivative of $f(y, z)$ with respect to the direction $\left(\frac{1}{\sqrt{2}}, \frac{-1}{\sqrt{2}}\right)$ at the point $(2, 1)$.

62.- It is known that the supply of a good as a function of his price p is given by:

$$S(p) = \begin{cases} \frac{p^2}{20} & \text{if } 0 \leq p < 10 \\ 2p - 15 & \text{if } 10 \leq p \leq 30 \end{cases}$$

- Determine the domain of the supply function.
- Which is the supply if the price is 5 monetary units?

63.- The demand of a good as a function of his price is given by $D(p) = ap^2 - b$. Determine the values of a and b knowing that $D(8) = 1$ and that $\lim_{p \rightarrow 0} D(p) = 5$.

64.- The total cost to produce q units of an article is $C(q) = 4q^3 - 6q^2 + 18q + 12$.

- Calculate the cost of producing 1 unit.
- If they are produced 20 units, which is the mean cost?
- Write the function of mean cost.
- Write the function of marginal cost.

65.- The profit in terms of the sale price of a product is $B(p)$, defined for $p \in [2, 6]$. This function $B(p)$ consist of two straight segments whose slopes are 0.2. Moreover it is known that $B(3)=50$ and that when the price is increased beyond the threshold of 4 monetary units a subsidy is lost so that the profit abruptly decreases by 10 units. Write and sketch graphically this profit function in terms of the price.

66.- The cost function in terms of the number of hours worked, x , is of the form $C(x) = \begin{cases} ax^2 + b & \text{if } x \in [0, 100] \\ c\sqrt{x} & \text{if } x \in (100, 200] \end{cases}$. Determine a , b , c knowing that $C(x)$ is continuous,

that the slope of the straight tangent at $x = 50$ is 8 and that 121 hours worked imply a cost equal to 990.

67.- The production function of a good is $Q(K, L) = 8\sqrt[3]{K^2L}$, where Q is the quantity of production, K and L are the quantity of inputs capital and labor, respectively.

Show that the productivity of the work is a function of the ratio capital-labor (only depends of the proportion between the capital and the labor).

68.- Let the production function of a firm be $Q(K, L) = 2K^{\frac{1}{2}}L^{\frac{2}{5}}$, where K is the capital, L the labour and Q the obtained production.

- a) Assuming that they are used 9 units of capital and 32 of work, what input increase implies greater production increase, keeping constant the other input?
- b) Assuming that they are used 9 units of capital and 32 of work, what would be the approximate variation in the production when the capital is increased 2 units and the labour 1 unit?
- c) Without computing the derivatives, calculate $K \frac{\partial Q}{\partial K}(K, L) + L \frac{\partial Q}{\partial L}(K, L)$.

69.- A firm produces two products with quantities q_1 y q_2 respectively, and the income is $I(q_1, q_2) = q_1 q_2$. Each one of the products has a production function that depends on the capital K and on the labor L as follows:

$$q_1(K, L) = 3K + 2L, \quad q_2(K, L) = 6\sqrt{K^2 L}$$

- a) Calculate $\frac{\partial I}{\partial K}(K, L), \frac{\partial I}{\partial L}(K, L)$ when they are used 4 units of capital and 9 of labor.
- b) Are the production functions homogeneous? of which degree?
- c) Without calculating the derivatives, compute $K \frac{\partial I}{\partial K}(K, L) + L \frac{\partial I}{\partial L}(K, L)$.

INTEGRALS OF FUNCTIONS OF ONE VARIABLE

1.- Compute the following indefinite integrals:

1) $\int \frac{3}{1+2y} dy$

2) $\int \frac{6z}{(z^2-5)^5} dz$

3) $\int \frac{1}{t^2} \sqrt{-1 + \frac{1}{t}} dt$

4) $\int (5 - 2x + \sqrt[4]{x^3} + 8e^x) dx$

5) $\int \frac{t}{1+t^2} dt$

6) $\int \frac{e^{\sqrt{y}}}{\sqrt{y}} dy$

7) $\int \left(\frac{1}{x} + \frac{1}{x^3} + 8 \frac{1}{\sqrt{x^3}} \right) dx$

8) $\int \left(\frac{1}{3}x + 5 \right)^9 dx$

9) $\int \frac{\ln t}{t} dt$

10) $\int \frac{5^t - 3^t}{7^t} dt$

11) $\int \frac{(\sqrt{x} - 2x)^2}{\sqrt{x}} dx$

12) $\int \frac{x-2}{x^2-4x+13} dx$

13) $\int \frac{(x+1)(x-1)}{x^2} dx$

14) $\int tg(x) dx$

15) $\int \sqrt[3]{\cos^2 x} \sin x dx$

16) $\int \frac{x}{1+x^2} dx$

17) $\int \frac{dx}{\sqrt{x}(\sqrt{x}+1)}$

18) $\int \frac{e^{2x}-1}{e^x-1} dx$

19) $\int \frac{2x+1}{x^2+x+1} dx$

20) $\int \frac{4-2x}{x^2-4x+3} dx$

21) $\int \frac{x-3}{x^2-6x+3} dx$

22) $\int x\sqrt{1-x^2} dx$

23) $\int \frac{x^2 dx}{(x^3-1)^2}$

24) $\int \frac{dx}{\sqrt{x}e^{\sqrt{x}}}$

2.- Compute the following definite integrals:

1) $\int_1^2 \left(x^3 - \frac{1}{x^2} + 2 \right) dx$

2) $\int_0^1 e^{-t+1} dt$

3) $\int_{-1}^1 \frac{x}{x^2+1} dx$

4) $\int_1^4 \left(\frac{1}{\sqrt{x^3}} - \sqrt{x} \right) dx$

5) $\int_{-2}^0 \left(\frac{x+4}{3} \right)^2 dx$

6) $\int_e^{e^2} \frac{1}{x \ln x} dx$

3.- Solve the following problems of application of integral calculus to the determination of areas:

1) Calculate the area limited by the graphics of the functions $3y = x^2$ and $y = -x^2 + 4x$.

2) Calculate the area of the figure limited by the parabolas $y = x^2 - 2x$ and $y = -x^2 + 4x$.

- 3) Calculate the area of the figure limited by the curves $y = x^2$, $y = 2x$ and $y = \frac{x^2}{2}$.
- 4) Calculate the area of the figure limited by the curves $x = 0$, $x = 2$, $y = 2^x$ and $y = 2x - x^2$.
- 5) Calculate the area limited by the curve $y = x^2 - 5x + 6$ and the line $y = 2x$.
- 6) Calculate the area limited by the parabola $y^2 = 4x$ and the straight line $y = x$.
- 7) Calculate the area limited by the parabola $y^2 = 4x$ and the line $y = 2x - 4$.
- 8) Calculate the area limited by the parabolas $x = -2y^2$ and $x = 1 - 3y^2$.

4.- Solve the following problems of application of integral calculus:

- 1) For a certain country, the marginal propensity to the consumption is given by $\frac{dC}{dI} = \frac{3}{4} - \frac{1}{2\sqrt{3I}}$, where the consumption C is a function of the national income I (in million euros). Determine the consumption function for that country if it is known that the consumption is 1 million euros when $I=12$.

- 2) The marginal cost function for a certain product is given by: $\frac{dc}{dq} = 10 - \frac{100}{q+10}$ where c is the total cost in euros when they are produced q units. When they are produced 100 units the mean cost is 50 euros each unit. Determine the fixed cost of the manufactured rounding off to the nearest integer of euro.

- 3) The marginal cost of the manufacture of a certain product is given by the function $CMg(x) = 6 - \frac{2}{\sqrt{x}}$, where x are the units produced. Knowing that the cost of operation is 84000 euros, find the total cost function of the manufacture of the cited product.

- 4) In the process of recovery of a patient, it is observed that the rhythm by which eliminates a toxic substance is given by the function $f(t) = -1.19e^{-0.22t}$, where t is the elapsed time in hours from the ingestion of the cited substance. Find the function that expresses the concentration of the substance in the blood, knowing that an hour after his ingestion the concentration is 1 gram by liter.

- 5) It is estimated that by t months the population of some city will change at a rate of $4 + 5t^{2/3}$ people by month. If the current population is 10.000 people, which is the population after 8 months have passed?

- 6) The promoters of a fair estimate that t hours after opening the doors, from 9:00 a.m., the visitors will enter to the fair at a rate of $N'(t)$ people by hour. Find an expression for the number of people that will go into the fair between the 11:00 a.m. and 1:00 p.m.
- 7) It is estimated that after t years have elapsed, the population of some community living on a lake shore will change at a rate of $0.6t^2 + 0.2t + 0.5$ thousands of people per year. The environmentalists have found that the level of pollution of the lake increases at a rate of, roughly, 5 units by 1.000 people. Find how much it will increase the pollution of the lake in the next 2 years.
- 8) A manufacturer claims that his marginal cost is $6q + 1$ euros by unit when they are produced q units. The total cost (included the indirect expenses) of producing the first unit is 130 euros. Which is the total cost of producing the first 10 units?